

Open camera or QR reader and
scan code to access this article
and other resources online.



Development of a Novel Anti-human EphA1 Monoclonal Antibody, Ea₁Mab-30, for Multiple Applications

Tomohiro Tanaka, Hiroyuki Suzuki, Honoka Taruta, Ayano Saga, Guanjie Li, Shiori Fujisawa, Mika K. Kaneko, and Yukinari Kato

Erythropoietin-producing hepatocellular receptor A1 (EphA1) is one of the Eph receptor family members, the largest group of receptor tyrosine kinases. EphA1 is expressed in various tissues and regulates cellular homeostasis by interacting with its membrane-bound ephrin ligands and other receptors. EphA1 critically correlates with the pathogenesis in several disorders, including Alzheimer's disease and cancers. Therefore, establishing sensitive monoclonal antibodies (mAbs) for EphA1 has been desired for basic research, diagnosis, and treatment. In this study, a novel specific and sensitive anti-human EphA1 mAb, clone Ea₁Mab-30 (mouse IgG₁, kappa), was established by the Cell-Based Immunization and Screening (CBIS) method. Ea₁Mab-30 demonstrated reactivity with an EphA1-overexpressed Chinese hamster ovary-K1 cell line (CHO/EphA1), an endogenously EphA1-expressing bladder carcinoma cell line (5637), and a colorectal adenocarcinoma cell line (Caco-2) in flow cytometry. Crossreactivities of Ea₁Mab-30 with other Eph receptors were not observed. Furthermore, the values of apparent binding affinity for CHO/EphA1 and 5637 were determined to be 8.9×10^{-9} M and 1.7×10^{-9} M, respectively. Furthermore, Ea₁Mab-30 detected EphA1 protein in CHO/EphA1 and 5637 lysates using Western blot analysis. Ea₁Mab-30 also clearly stained EphA1 of formalin-fixed paraffin-embedded CHO/EphA1 using immunohistochemistry. Ea₁Mab-30, established by CBIS method, could help analyze the EphA1-contributed cellular functions and have potential applications in pathological diagnosis and treatment with specificity and high affinity for EphA1-expressing cells.

Keywords: Eph receptor, EphA1, CBIS method, monoclonal antibody, flow cytometry, Western blot, immunohistochemistry

Introduction

Erythropoietin-producing hepatocellular (Eph) receptors are single-pass transmembrane glycoproteins, which belong to the most prominent family of receptor tyrosine kinases.^{1,2} Eph receptors play pivotal roles in tissue homeostasis, including cell proliferation, migration, and angiogenesis by binding to their ligands, ephrin.^{1,2} Since it was first identified in hepatocellular carcinoma cell line in 1987, 14 Eph receptors (EphA1 to EphA8, EphA10, EphB1 to EphB4, and EphB6) and eight ephrin ligands [glycosylphosphatidylinositol-anchored ephrin A1 to A5 and transmembrane ephrin B1 to B3] have been identified so far.³ Eph receptors are topics of potential targets for diseases such as cancers because the

aberrant functions of Eph receptors disrupt cellular homeostasis and induce tumor development.

EphA1 was first isolated from an erythropoietin-producing human hepato-cellular carcinoma cell line (ETL-1).³ EphA1 is expressed in epithelial and nervous tissues, including skin, neuron, and colon crypt base, and controls many cellular functions, such as cell morphology, motility, axonal guidance, and synaptic plasticity.⁴⁻⁸ EphA1 expression is also confirmed in T-lymphocytes, monocytes.⁹ EphA1-knockout (KO) mice exhibited abnormal tail development as a kinky-tail, and an uterovaginal defection caused by a deficiency in apoptosis during reproductive tract canalization.¹⁰ EphA1 plays an essential role in normal tissue development. According to the genome-wide association studies (GWAS) of Alzheimer's

disease (AD) patients, EphA1 has been considered as an associated factor for AD.^{11–13} AD is a progressive age-associated neurodegenerative disorder leading to loss of autonomy, which pathologically defined by extensive neuronal loss, the accumulation of neurofibrillary tangles, and extracellular amyloid plaques in the brain.^{14–16} Missense mutations in EphA1 gene have also been reported that may contribute to the development of AD by disrupting receptor signaling.¹⁷ Some single nucleotide polymorphisms (SNPs), rs11767557 and rs11771145, which affect the EphA1 gene, have been reported to decrease the risk of late-onset AD (patients of over 65 years old).^{11,13,18} Furthermore, rs11767557 variant is associated with EphA1 transcription in human whole blood data analysis. Meanwhile, rs11767557 SNP has not shown the significant effect of EphA1 transcription in brain tissue.¹⁹ Moreover, EphA1 transcription is not altered in brain with AD.²⁰ The discrepancies in these results may be due to the size and heterogeneity of the samples. Since few molecular analyses have been conducted to clarify the regulation of EphA1 in AD, further investigation is required to determine whether EphA1 is involved in the risk of AD.

The EphA1 upregulation is also correlated in various cancers, including esophageal,²¹ colon,³ gastric,^{22,23} and prostate cancers.^{24,25} EphA1 high expression associates with unfavorable overall survival and higher tumor stage.^{21,26} EphA1 has been implicated as a marker for aggressive ovarian cancer stem cells with coexpression with aldehyde dehydrogenase 1A1, CD9, and CD24.²⁷ Furthermore, a disintegrin and metalloproteinase 12 (ADAM12)-cleaved ephrinA1 facilitates tumor cell entry into the lungs by activating EphA1.²⁸ While EphA1 is focused as a cancer-promoting factor, tumor suppressor functions of EphA1 have also been proposed. Reduced expression of EphA1 correlates with tumor invasion, lymph node metastasis, and poor outcomes in several cancers, including colon, skin, and ovarian cancers.^{29–33} The inhibition of cell spreading and Rac1 activation, which promotes cell motility, is observed by EphA1 activation.⁵ It is necessary to clarify the role of EphA1 in promoting or suppressing tumors.

Previously, we have successfully developed several monoclonal antibodies (mAbs) against Eph receptors, including human EphA2 (clone Ea₂Mab-7),³⁴ human EphB2 (clone Eb₂Mab-12),³⁵ human EphB4 (clone B4Mab-7),³⁶ and human EphB6 (clone Eb₆Mab-3)³⁷ by using the Cell-Based Immunization and Screening (CBIS) method. The CBIS method is an efficient strategy for developing antibodies with a wide variety of epitopes such as structural epitopes and linear epitopes, as well as modifications of cell surface membrane proteins in a short period. In this study, we have established a novel anti-human EphA1 mAb (Ea₁Mab-30) using the CBIS method.

Materials and Methods

Preparation of cell lines

Chinese hamster ovary (CHO)-K1 (CCL-61), LN229 glioblastoma (CRL-2611), P3X63Ag8U.1 (P3U1, CRL-1597) myeloma, and Caco-2 (HTB-37) colorectal adenocarcinoma were obtained from the American Type Culture Collection (Manassas, VA, USA). 5637 bladder carcinoma (TKG 0605) was obtained from the Cell Resource Center for Biomedical Research, Institute of Development, Aging and Cancer,

Tohoku University (Miyagi, Japan). The complementary DNA (cDNA) of human EphA1 [Catalog No.: RC213689 (OriGene Technologies, Inc., Rockville, MD, USA), Accession No.: NM_005232] plus an N-terminal MAP tag (GDGMVPPGIEDK)³⁸ recognized by an anti-MAP tag mAb (PMab-1) or an N-terminal PA tag (GVAMPGAEDDVV)³⁹ recognized by an anti-PA tag mAb (NZ-1), were subcloned into a pCAGzeo vector [FUJIFILM Wako Pure Chemical Corporation (Wako), Osaka, Japan]. Afterward, plasmids were transfected into CHO-K1 and LN229 cells and stable transfectants [CHO/MAP-EphA1, CHO/PA-EphA1 (CHO/EphA1), LN229/MAP-EphA1, and LN229/PA-EphA1 (LN229/EphA1)] were subsequently selected using a cell sorter (SH800, Sony Corp., Tokyo, Japan). After sorting, cultivation in a medium containing 0.5 mg/mL of Zeocin (InvivoGen, San Diego, CA, USA) was conducted.

GeneArt™ CRISPR nuclease vectors with OFP plasmid which target human EphA1 (TrueGuide™ Synthetic sgRNA, CRISPR466292_SGM) was purchased from Thermo Fisher Scientific Inc. The EphA1 KO 5637 cell line (BINDS-38) was generated by transfecting the vector using the Neon Transfection System (Thermo Fisher Scientific, Inc.). Stable transfectants were established by cell sorting using SH800 (Sony Corp.).

Other Eph receptor-expressing CHO-K1 cells (e.g., CHO/EphA2) were established and cultured as previously reported.³⁵

Antibodies

A purified anti-human EphA1 Antibody (clone A20047D, rat IgG_{2a}, kappa) was purchased from BioLegend (San Diego, CA, USA). An anti-isocitrate dehydrogenase 1 (IDH1) mAb (clone RcMab-1) was developed previously.⁴⁰ Alexa Fluor 488-conjugated anti-mouse IgG and anti-rat IgG were purchased from Cell Signaling Technology, Inc. (Danvers, MA, USA). Horseradish peroxidase-conjugated anti-mouse IgG and anti-rat IgG were obtained from Agilent Technologies Inc. (Santa Clara, CA, USA) and Merck KGaA (Darmstadt, Germany), respectively.

Production of hybridomas

Two 5-week-old female BALB/cAJcl mice were purchased from CLEA Japan (Tokyo, Japan). The animal experiment was approved by the Animal Care and Use Committee of Tohoku University (Permit number: 2019NiA-001). For developing anti-human EphA1 mAbs, the immunogen CHO/MAP-EphA1 cells were harvested after brief treatment with 1 mM ethylenediaminetetraacetic acid (EDTA; Nacalai Tesque, Inc.) to avoid antigen digestion.

In the first immunization, the CHO/MAP-EphA1 (1×10^8 cells/mouse) mixed with Inject Alum Adjuvant (Thermo Fisher Scientific Inc.) was immunized via the intraperitoneal route. Three additional injections of CHO/MAP-EphA1 (1×10^8 cells/mouse) without the adjuvant were conducted intraperitoneally in every week. A last booster injection was performed with 1×10^8 cells/mouse of CHO/MAP-EphA1 via intraperitoneal route two days before harvesting spleen cells from mice. We gently executed cell-fusion of P3U1 myeloma cells with the harvested splenocytes from immunized mice using polyethylene glycol 1500 (PEG1500; Roche Diagnostics, Indianapolis, IN, USA) at 37°C.

Hybridomas were cultured in the RPMI-1640 medium added some supplements as shown above, with additional supplements included hypoxanthine, aminopterin, and thymidine (HAT; Thermo Fisher Scientific, Inc.), 5% BriClone (NICB, Dublin, Ireland), and 5 µg/mL of Plasmocin (InvivoGen) as antibiotics into the medium. On days 6 to 7 after cell-fusion, the hybridoma supernatants were screened by flow cytometry using LN229/MAP-EphA1 and parental

LN229 cells. The hybridoma supernatant containing Ea₁Mab-30 in serum-free medium was filtered and purified using Ab-Capcher Extra (ProteNova, Kagawa, Japan).

Flow cytometric analysis

Cells were harvested using 1 mM EDTA (Nacalai Tesque, Inc.). Subsequently, cells were gently washed with 0.1%




FIG. 1. A schematic depiction of anti-human EphA1 mAbs development by CBIS method. **(A)** CHO/MAP-EphA1 cells were immunized into two female mice via the intraperitoneal route. **(B)** The spleen cells isolated from antigen-immunized mice were fused with P3U1. **(C)** The culture supernatants of hybridoma were screened by flow cytometric analysis using LN229 and LN229/MAP-EphA1 to select EphA1-specific mAb-producing hybridomas. **(D)** Single hybridoma clones were obtained by limiting dilution, followed by additional screening. Finally, Ea₁Mab-30 (mouse IgG₁, kappa) was established. CBIS, Cell-Based Immunization and Screening; Ea₁Mab-30, anti-human EphA1 mAb; EphA1, erythropoietin-producing hepatocellular receptor A1; mAb, monoclonal antibody.

bovine serum albumin in phosphate-buffered saline (PBS) and treated with primary mAbs for 30 minutes at 4°C. After that, cells were treated with Alexa Fluor 488-conjugated anti-mouse or rat IgG (dilution rate; 1:1000) following the fluorescence data collection, using the SA3800 Cell Analyzer (Sony Corp.).

To determine the isotype of mAbs, primary mAb-treated CHO/EphA1 cells were incubated with Alexa Fluor 488-conjugated anti-mouse immunoglobulins, fluorescein isothiocyanate (FITC)-conjugated anti-mouse heavy chains (IgG₁, IgG_{2a}, IgG_{2b}, IgG₃, and IgM; SouthernBiotech, Birmingham, AL, USA), or FITC-conjugated anti-mouse light chains (kappa

and lambda; SouthernBiotech). The fluorescence data were analyzed using the SA3800 Cell Analyzer.

Determination of the apparent binding affinity by flow cytometry

CHO/EphA1 and 5637 cells were suspended in 100 μ L serially diluted of Ea₁Mab-30 (20 μ g/mL to 0.0012 μ g/mL) and A20047D (25 μ g/mL to 0.0015 μ g/mL), after which Alexa Fluor 488-conjugated anti-mouse or rat IgG (dilution rate; 1:200) was treated. Fluorescence data were subsequently




FIG. 2. Flow cytometric analysis of anti-human EphA1 mAbs against exogenously expressed human EphA1. CHO/EphA1 (A) and CHO-K1 (B) cells were treated with 0.01–10 μ g/mL of Ea₁Mab-30 or A20047D (red line), followed by treatment with Alexa Fluor 488-conjugated anti-mouse or rat IgG. Fluorescence data were collected using the SA3800 Cell Analyzer. Black line, control (no primary antibody treatment). A20047D, anti-human EphA1 antibody.

collected, using the BD FACSLyric (BD Biosciences, Franklin Lakes, NJ, USA), following the analyzing of the apparent binding affinity (K_D) as described previously.³⁵

Western blot analysis

Cells were lysed and boiled in sodium dodecyl sulfate sample buffer (Nacalai Tesque, Inc.). Denatured proteins were electrophoresed and transferred onto polyvinylidene difluoride membranes (Merck KGaA). After blocking with 4% non-fat milk (Nacalai Tesque, Inc.), the membranes were treated with Ea₁Mab-30 (1 µg/mL), A20047D (1 µg/mL), an anti-IDH1 mAb (clone RcMab-1, 1 µg/mL), or an anti-PA tag mAb (clone NZ-1, 1 µg/mL), followed by incubation with horseradish peroxidase-conjugated anti-mouse IgG for Ea₁Mab-30 (dilution rate; 1:2000; Agilent Technologies Inc.) or anti-rat IgG for A20047D, RcMab-1, and NZ-1 (dilution rate; 1:10000; Merck KGaA). Chemiluminescence signals were visualized using ImmunoStar LD (Wako) and imaged with a Sayaca-Imager (DRC Co. Ltd., Tokyo, Japan).

Immunohistochemical analysis

The formalin-fixed paraffin-embedded (FFPE) CHO/EphA1 and CHO-K1 cell blocks were prepared using iPGell (Genostaff Co., Ltd., Tokyo, Japan). The antigen retrieval of paraffin-embedded cell sections was conducted by autoclave in a citrate buffer (pH 6.0; Nichirei Biosciences, Inc., Tokyo, Japan). After blocking using the SuperBlock T20 Blocking Buffer (Thermo Fisher Scientific Inc.), the sections were treated with Ea₁Mab-30 (5 µg/mL) and A20047D (5 µg/mL) and then reacted with the Envision+ Kit (Agilent Technologies Inc.) for Ea₁Mab-30 or Histofine Simple Stain Mouse MAX PO (Rat) (Nichirei Biosciences, Inc.) for A20047D. Color development and counterstaining were performed as described previously.³⁶

Results

Development of anti-EphA1 mAbs using the CBIS method

To establish mAbs targeting human EphA1, we employed the CBIS method using human EphA1-overexpressed cell lines. Anti-human EphA1 mAbs-producing hybridoma were screened by flow cytometric analysis (Fig. 1). Two female BALB/cAJcl mice were immunized with CHO/MAP-EphA1 (1×10^8 cells/time/mouse) every week via intraperitoneal route for a total of 5 times. Subsequently, splenocytes isolated from immunized mice and P3U1 cells were fused by PEG1500 and fused cells were seeded into 96-well plates. After confirming hybridoma formation, flow cytometric high throughput screening was conducted to select LN229/MAP-EphA1-reactive and parental LN229-nonreactive supernatants of hybridomas. Afterward, we obtained highly LN229/MAP-EphA1-reactive supernatants of the hybridoma. After limiting dilution of each positive well, we investigated the reactivity of several anti-EphA1 mAbs to CHO/EphA1 in flow cytometry

and Western blot analysis. Finally, we selected the highly sensitive clone Ea₁Mab-30 (mouse IgG₁, kappa). We also established more clones with Ea₁Mab-30. The information has been updated on our website "Antibody bank" (http://www.med-tohoku-antibody.com/topics/001_paper_antibody_PDIS.htm#EphA1).

Investigation of antibody reactivities using flow cytometry

We checked the reactivity of Ea₁Mab-30 against CHO-K1 and CHO/EphA1 cells by flow cytometric analysis. A commercially available anti-human EphA1 mAb from BioLegend, A20047D, was used for comparison with Ea₁Mab-30. A20047D is currently the only commercially available mAb that can be used for flow cytometry. Results showed that Ea₁Mab-30 and A20047D recognized CHO/EphA1 dose-dependently (Fig. 2A). Reactivity is almost the same for Ea₁Mab-30 and A20047D to CHO/EphA1 (Fig. 2A). Neither Ea₁Mab-30 nor A20047D reacted with parental CHO-K1 cells (Fig. 2B). Thus, Ea₁Mab-30 can detect exogenously expressed human EphA1 specifically in flow cytometry.

The reactivity of anti-EphA1 mAbs using flow cytometry against endogenous EphA1


Next, we confirmed the reactivity of Ea₁Mab-30 and A20047D in flow cytometry against 5637 and Caco-2 cells which are endogenously human EphA1 expressing cells. The high expression of Eph receptor and ephrin have been reported in bladder cancers.⁴¹ We initially confirmed that EphA1 was endogenously expressed in the bladder cancer cell line, 5637. Furthermore, the expression of EphA1 in colorectal cancer, including Caco-2 cells, has been described in previous reports.^{30,42} Results showed that Ea₁Mab-30 and A20047D recognized endogenously expressing EphA1 in 5637 (Fig. 3A) and Caco-2 (Fig. 3B) cancer cells. Ea₁Mab-30 showed slightly higher reactivity than A20047D on all concentrations (Fig. 3A and 3B). Neither Ea₁Mab-30 nor A20047D reacted with BINDS-38, EphA1-KO 5637 cells, even at a concentration of 10 µg/mL (Fig. 3C). Thus, Ea₁Mab-30 can detect endogenously expressing EphA1 specifically in flow cytometry.

Specificity of Ea₁Mab-30 to Eph receptor-overexpressed CHO-K1 cells


We have established the cell lines of all Eph receptor-overexpressed CHO-K1 cells, EphA1 to A8, A10, B1 to B4, and B6, respectively. Using the 14 cell lines, the specificity of Ea₁Mab-30 was analyzed. As shown in Figure 4, 10 µg/mL of Ea₁Mab-30 potently recognized CHO/EphA1. Ea₁Mab-30 never reacted to other Eph receptors. This result complemented the fact that Ea₁Mab-30 is an EphA1-specific mAb.

FIG. 3. Flow cytometric analysis of anti-human EphA1 mAbs against endogenously expressing human EphA1 in cancer cells. 5637 (A), Caco-2 (B), and BINDS-38 (C) cells were treated with 0.01–10 µg/mL of Ea₁Mab-30 or A20047D (red line), followed by treatment with Alexa Fluor 488-conjugated anti-mouse or rat IgG. Fluorescence data were collected using the SA3800 Cell Analyzer. Black line, control (no primary antibody treatment). Caco-2, colorectal adenocarcinoma cell line.


A



B



C



Calculation of the apparent binding affinity of anti-EphA1 mAbs using flow cytometry

The apparent binding affinity of Ea₁Mab-30 and A20047D was assessed with exogenously EphA1-expressed CHO/EphA1 and endogenously EphA1-expressing 5637 using flow cytometry. The results indicated that the K_D values of Ea₁Mab-30 and A20047D for CHO/EphA1 were 8.9×10^{-9} M and 8.6×10^{-9} M, respectively (Fig. 5A). The K_D values of Ea₁Mab-30 and A20047D for 5637 were 1.7×10^{-9} M and 2.7×10^{-9} M, respectively (Fig. 5B). Although there was no noticeable difference in binding affinity for CHO/EphA1 between Ea₁Mab-30 and A20047D, Ea₁Mab-30 possessed higher binding affinity for 5637 than A20047D. These results demonstrate that Ea₁Mab-30 can recognize EphA1 with high affinity to EphA1 on cells.

Western blot analysis using anti-EphA1 mAbs

We investigated whether Ea₁Mab-30 can be used for Western blot analysis by analyzing LN229, EphA1-overexpressed LN229/EphA1, 5637, and EphA1-KO BINDS-38 cell lysates. The estimated molecular weight of EphA1 protein is 108 kDa. As shown in Figure 6A, Ea₁Mab-30 could detect exogenously expressed-EphA1 as the band around 100 to 130 kDa in LN229/EphA1 cell lysates, while no band was detected in parental LN229 cells. Another anti-EphA1 mAb A20047D could not detect any band in LN229/EphA1 cell lysates. Western blot has not been confirmed to apply in the supplier data source from BioLegend. An anti-PA tag mAb NZ-1 was used as a positive control and could also detect a band of the same position around 100 to 130 kDa in LN229/EphA1 cell lysates. An anti-IDH1 mAb (clone RcMab-1)




FIG. 4. Crossreactivity of Ea₁Mab-30 in Eph receptor-expressed CHO-K1 cells. CHO-K1 cells, which overexpressed each of the 14 Eph receptors, were treated with 10 µg/mL of Ea₁Mab-30 (red line) or control blocking buffer (black line) followed by the treatment with anti-mouse IgG conjugated with Alexa Fluor 488. Fluorescence data were collected using the SA3800 Cell Analyzer. Black line, control (no primary antibody treatment).




FIG. 5. Evaluation of the K_D values of anti-human EphA1 mAbs. **(A)** CHO/EphA1 cells were suspended in 100 μL of serially diluted 20 $\mu\text{g/mL}$ to 0.0012 $\mu\text{g/mL}$ of Ea₁Mab-30 and 20 $\mu\text{g/mL}$ to 0.0012 $\mu\text{g/mL}$ of A20047D. **(B)** 5637 cells were suspended in 100 μL of serially diluted 25 $\mu\text{g/mL}$ to 0.0015 $\mu\text{g/mL}$ of Ea₁Mab-30 and 25 $\mu\text{g/mL}$ to 0.0015 $\mu\text{g/mL}$ of A20047D. After treatments of primary mAbs, cells were treated with Alexa Fluor 488-conjugated anti-mouse IgG (for Ea₁Mab-30) or anti-rat IgG (for A20047D). Subsequently, the K_D values were determined using the geometric mean.

was used for internal control. Furthermore, Ea₁Mab-30 could detect endogenously expressing-EphA1 as the band around 100 to 130 kDa in 5637 cell lysates, while no band was detected in BINDS-38, EphA1-KO 5637 cells (Fig. 6B). A20047D could not show any band in 5637 cell lysates. These results demonstrate that in Western blot analysis, Ea₁Mab-30 can detect EphA1 in exogenously and endogenously EphA1-expressing cells.

Immunohistochemistry using anti-EphA1 mAbs

To evaluate whether Ea₁Mab-30 can be used for immunohistochemistry (IHC), the FFPE sections of CHO-K1 and CHO/EphA1 were stained with Ea₁Mab-30. Apparent membranous staining by Ea₁Mab-30 was observed in CHO/EphA1 (Fig. 7A). Another anti-EphA1 mAb, A20047D did not stain CHO/EphA1 sections (Fig. 7A). Both Ea₁Mab-30 and A20047D did not react with the parental CHO-K1 sections

(Fig. 7B). These results indicate that Ea₁Mab-30 applies to IHC for detecting EphA1-positive cells in paraffin-embedded cell samples.

Discussion

Currently, no mAbs targeting human EphA1 can be used for flow cytometry, Western blot, and immunohistochemistry. We successfully established a novel anti-human EphA1 mAb, Ea₁Mab-30, which is applicable for various experiments. Ea₁Mab-30 specifically detected EphA1 in flow cytometry (Figs 2 to 4), Western blot (Fig. 6), and immunohistochemistry (Fig. 7). Furthermore, Ea₁Mab-30 recognized EphA1 with higher affinity than A20047D, the only commercially available mAb suitable for flow cytometry (Fig. 5). Since the estimation of K_D by flow cytometry lacks the precision and kinetics, further analyses such as surface plasmon resonance or biolayer interferometry are




FIG. 6. Detection of EphA1 protein using Ea₁Mab-30 by Western blot analysis. Cell lysates (10 μ g/lane) of LN229, LN229/EphA1 (**A**), 5637, and BINDS-38 (**B**) were electrophoresed on polyacrylamide gels and transferred onto polyvinylidene difluoride membranes. The membranes were incubated with Ea₁Mab-30 (1 μ g/mL), A20047D (1 μ g/mL), an anti-PA tag mAb (clone NZ-1, 1 μ g/mL), and RcMab-1 (1 μ g/mL) and subsequently with horseradish peroxidase-conjugated anti-mouse or anti-rat immunoglobulins. RcMab-1, anti-isocitrate dehydrogenase 1 mAb.

essential. Nevertheless, Ea₁Mab-30 could contribute to elucidating the EphA1-related cellular functions and diagnosis of disorders such as AD and tumors. Overexpression of EphA1 is observed in various cancers,^{23,33,43} EphA1 can be a target for cancer treatments. To evaluate the anti-tumor efficacy of Ea₁Mab-30, it will be converted to a mouse IgG_{2a} version to elevate antibody-dependent cellular cytotoxicity in the next step.^{44,45}

Since overexpression of Eph receptors has been observed regardless of cancer type, they have been widely studied in cancer research and emerged as therapeutic targets.^{46–49} However, specific drugs targeting Eph receptors or ephrin ligands have yet to be approved. However, some clinical trials have been conducted such as DS-8895a (a humanized anti-EphA2 afucosylated mAb) against advanced EphA2-expressing cancer and Ifabotuzumab (an anti-EphA3 mAb)




FIG. 7. Immunohistochemical staining of paraffin-embedded sections of CHO/EphA1 and CHO-K1. The sections of CHO/EphA1 (**A**) and CHO-K1 (**B**) cells were treated with Ea₁Mab-30 (5 μg/mL) or A20047D (5 μg/mL), followed by that with the Envision + Kit (for Ea₁Mab-30) or Histofine Simple Stain Mouse MAX PO (Rat) (for A20047D). Color was developed using 3,3'-diaminobenzidine tetrahydrochloride, and counterstaining was performed using hematoxylin. Scale bar = 100 μm.

against advanced hematological malignancies.^{48,50} EphA1-targeting therapeutic challenge has not been confirmed yet. One reason could be that EphA1 has a dual role in promoting and suppressing cancer. Since Ea₁Mab-30 can detect EphA1 in immunohistochemistry using the FFPE sections of CHO/EphA1, further studies are essential whether Ea₁Mab-30 can detect EphA1 in the FFPE sections of various tumors for diagnosis.

AD is the most common chronic neurodegenerative disease characterized by cognitive decline and behavioral change. The most well-known neuropathological hallmarks of AD are aggregation of extracellular amyloid- β peptides into amyloid plaques, and cytotoxic intracellular neurofibrillary tau tangles.^{51,52} Large number of GWAS for AD have been conducted and suggested numerous loci as the risk of AD including EphA1.^{13,14,18,53} Some SNPs near EphA1 gene, rs11767557 and rs11771145, have been demonstrated in GWAS to decrease the risk of AD.^{13,18,54} Several EphA1 missense mutations are considered to increase the risk of AD. P460L mutation in EphA1 is associated with AD by dysregulating EphA1 activity.⁵⁵ P460L variant triggers disruptions of EphA1, including reducing tyrosine phosphorylation, cell surface expression, and decreasing T-cell recruitment to the brain by affecting brain endothelial barrier.⁵⁵ In *Drosophila* models, P460L mutation causes some AD-like symptoms such as hyperexcitability, less sleep, a more vigorous circadian rhythm and overactivity of clock neuron in the absence of memory loss.⁵⁶ P460 locates in the second fibronectin type III repeat (FNIII-2) of EphA1 adjacent to the cell membrane and highly conserved across species.^{3,55,57} Thus, this missense mutation might be necessary for neuropathological research in animal models. EphA-ephrinA axis allows forward and reverse bidirectional

signaling in cell-cell contact, followed by playing a role in synaptic formation and plasticity, axonal guidance, and brain development.^{5,7,58} EphA1 is an attractive molecule for analyzing AD pathology, but there are few reports of postmortem brain-tissue analysis. Moreover, EphA1 promotes the pathogenesis of Parkinson's disease, the second most common neurodegenerative disease, through CXCL12-CXCR4 axis.⁵⁹ The developed Ea₁Mab-30 will be helpful for the analysis of the diseases.

Although EphA1 is characterized as the negative risk factor of AD, its structural alteration with missense mutations may provoke the AD.⁶⁰ As with cancer, to clarify the critical mutation of AD-related proteins is essential to make an early diagnosis and treatment. Furthermore, N414 and N478 in EphA1 are glycosylated.¹⁷ Although these *N*-glycosylation are impaired by P460L mutation, epigenetic modification might contribute to normal EphA1 functions. We have previously identified cancer specific aberrant glycosylation of podoplanin⁶¹ and misfolding of HER2⁴⁴ through the establishment of mAbs targeting these characteristic epitopes. Research into disease-specific epitopes for EphA1, such as those for cancer and AD, is expected to progress. We plan to obtain mutated-EphA1 specific mAbs to clarify the relation between EphA1 and diseases.

Authors' Contributions

T.T., H.S., H.T., A.S., G.L., and S.F. performed the experiments. M.K.K. and Y.K. designed the experiments. T.T. analyzed the data. T.T. wrote the article. All the authors have read and agreed to the published version of the article.

Author Disclosure Statement

No competing financial interests exist.

Funding Information

This research was supported in part by Japan Agency for Medical Research and Development (AMED) under Grant Numbers: JP25am0521010 (to Y.K.), JP25ama121008 (to Y.K.), JP25ama221339 (to Y.K.), and JP25bm1123027 (to Y.K.), and by the Japan Society for the Promotion of Science (JSPS) Grants-in-Aid for Scientific Research (KAKENHI) grant nos. 24K18268 (to T.T.) and 25K10553 (to Y.K.).

References

1. Wang HU, Chen ZF, Anderson DJ. Molecular distinction and angiogenic interaction between embryonic arteries and veins revealed by ephrin-B2 and its receptor Eph-B4. *Cell* 1998;93(5):741–753; doi: 10.1016/s0092-8674(00)81436-1
2. Sawamiphak S, Seidel S, Essmann CL, et al. Ephrin-B2 regulates VEGFR2 function in developmental and tumour angiogenesis. *Nature* 2010;465(7297):487–491; doi: 10.1038/nature08995
3. Hirai H, Maru Y, Hagiwara K, et al. A novel putative tyrosine kinase receptor encoded by the eph gene. *Science* 1987;238(4834):1717–1720; doi: 10.1126/science.2825356
4. Coulthard MG, Lickliter JD, Subanese N, et al. Characterization of the EphA1 receptor tyrosine kinase: Expression in epithelial tissues. *Growth Factors* 2001;18(4):303–317; doi: 10.3109/08977190109029118

5. Yamazaki T, Masuda J, Omori T, et al. EphA1 interacts with integrin-linked kinase and regulates cell morphology and motility. *J Cell Sci* 2009;122(Pt 2):243–255; doi: 10.1242/jcs.036467
6. Martínez A, Otal R, Sieber BA, et al. Disruption of ephrin-A/EphA binding alters synaptogenesis and neural connectivity in the hippocampus. *Neuroscience* 2005;135(2):451–461; doi: 10.1016/j.neuroscience.2005.06.052
7. Lai KO, Ip NY. Synapse development and plasticity: Roles of ephrin/Eph receptor signaling. *Curr Opin Neurobiol* 2009;19(3):275–283; doi: 10.1016/j.conb.2009.04.009
8. Kosinski C, Li VS, Chan AS, et al. Gene expression patterns of human colon tops and basal crypts and BMP antagonists as intestinal stem cell niche factors. *Proc Natl Acad Sci U S A* 2007;104(39):15418–15423; doi: 10.1073/pnas.0707210104
9. Sakamoto A, Sugamoto Y, Tokunaga Y, et al. Expression profiling of the ephrin (EFN) and Eph receptor (EPH) family of genes in atherosclerosis-related human cells. *J Int Med Res* 2011;39(2):522–527; doi: 10.1177/147323001103900220
10. Duffy SL, Coulthard MG, Spanevello MD, et al. Generation and characterization of EphA1 receptor tyrosine kinase reporter knockout mice. *Genesis* 2008;46(10):553–561; doi: 10.1002/dvg.20434
11. Lambert JC, Ibrahim-Verbaas CA, Harold D, et al.; Cohorts for Heart and Aging Research in Genomic Epidemiology. Meta-analysis of 74,046 individuals identifies 11 new susceptibility loci for Alzheimer's disease. *Nat Genet* 2013;45(12):1452–1458; doi: 10.1038/ng.2802
12. Talebi M, Delpak A, Khalaj-Kondori M, et al. ABCA7 and EphA1 genes polymorphisms in late-onset Alzheimer's disease. *J Mol Neurosci* 2020;70(2):167–173; doi: 10.1007/s12031-019-01420-x
13. Naj AC, Jun G, Beecham GW, et al. Common variants at MS4A4/MS4A6E, CD2AP, CD33 and EPHA1 are associated with late-onset Alzheimer's disease. *Nat Genet* 2011;43(5):436–441; doi: 10.1038/ng.801
14. Karch CM, Goate AM. Alzheimer's disease risk genes and mechanisms of disease pathogenesis. *Biol Psychiatry* 2015;77(1):43–51; doi: 10.1016/j.biopsych.2014.05.006
15. Sadigh-Eteghad S, Sabermarouf B, Majdi A, et al. Amyloid-beta: A crucial factor in Alzheimer's disease. *Med Princ Pract* 2015;24(1):1–10; doi: 10.1159/000369101
16. Hardy J, Selkoe DJ. The amyloid hypothesis of Alzheimer's disease: Progress and problems on the road to therapeutics. *Science* 2002;297(5580):353–356; doi: 10.1126/science.1072994
17. Matsumoto M, Gomez-Soler M, Lombardi S, et al. Missense mutations of the ephrin receptor EPHA1 associated with Alzheimer's disease disrupt receptor signaling functions. *J Biol Chem* 2025;301(2):108099; doi: 10.1016/j.jbc.2024.108099
18. Hollingworth P, Harold D, Sims R, et al.; EADI1 consortium. Common variants at ABCA7, MS4A6A/MS4A4E, EPHA1, CD33 and CD2AP are associated with Alzheimer's disease. *Nat Genet* 2011;43(5):429–435; doi: 10.1038/ng.803
19. Liu G, Zhang Y, Wang L, et al. Alzheimer's Disease rs11767557 Variant Regulates EPHA1 Gene Expression Specifically in Human Whole Blood. *J Alzheimers Dis* 2018;61(3):1077–1088; doi: 10.3233/jad-170468
20. Karch CM, Jeng AT, Nowotny P, et al. Expression of novel Alzheimer's disease risk genes in control and Alzheimer's disease brains. *PLoS One* 2012;7(11):e50976; doi: 10.1371/journal.pone.0050976
21. Wang J, Ma J, Dong Y, et al. High expression of EphA1 in esophageal squamous cell carcinoma is associated with lymph node metastasis and advanced disease. *APMIS* 2013;121(1):30–37; doi: 10.1111/j.1600-0463.2012.02941.x
22. Wang J, Dong Y, Wang X, et al. Expression of EphA1 in gastric carcinomas is associated with metastasis and survival. *Oncol Rep* 2010;24(6):1577–1584; doi: 10.3892/or.00001020
23. Wang YC, Dai Y, Xu GL, et al. Association Between EphA1 and Tumor Microenvironment in Gastric Carcinoma and its Clinical Significance. *Med Sci Monit* 2020;26:e923409; doi: 10.12659/msm.923409
24. Robinson D, He F, Pretlow T, et al. A tyrosine kinase profile of prostate carcinoma. *Proc Natl Acad Sci U S A* 1996;93(12):5958–5962; doi: 10.1073/pnas.93.12.5958
25. Peng L, Wang H, Dong Y, et al. Increased expression of EphA1 protein in prostate cancers correlates with high Gleason score. *Int J Clin Exp Pathol* 2013;6(9):1854–1860.
26. Koh HM, Jang BG, Lee DH, et al. The association between ephrin receptor-A1 expression and survival in patients with cancer: A meta-analysis. *Transl Cancer Res* 2022;11(6):1587–1594; doi: 10.21037/tcr-21-1367
27. Nagare RP, Sneha S, Krishnapriya S, et al. ALDH1A1+ ovarian cancer stem cells co-expressing surface markers CD24, EPHA1 and CD9 form tumours in vivo. *Exp Cell Res* 2020;392(1):112009; doi: 10.1016/j.yexcr.2020.112009
28. Ieguchi K, Tomita T, Omori T, et al. ADAM12-cleaved ephrin-A1 contributes to lung metastasis. *Oncogene* 2014;33(17):2179–2190; doi: 10.1038/onc.2013.180
29. Dong Y, Wang J, Sheng Z, et al. Downregulation of EphA1 in colorectal carcinomas correlates with invasion and metastasis. *Mod Pathol* 2009;22(1):151–160; doi: 10.1038/modpathol.2008.188
30. Herath NI, Doecke J, Spanevello MD, et al. Epigenetic silencing of EphA1 expression in colorectal cancer is correlated with poor survival. *Br J Cancer* 2009;100(7):1095–1102; doi: 10.1038/sj.bjc.6604970
31. Hafner C, Becker B, Landthaler M, et al. Expression profile of Eph receptors and ephrin ligands in human skin and downregulation of EphA1 in nonmelanoma skin cancer. *Mod Pathol* 2006;19(10):1369–1377; doi: 10.1038/modpathol.3800660
32. Jin Y, Zou Y, Wan L, et al. Decreased Eph receptor-A1 expression is related to grade in ovarian serous carcinoma. *Mol Med Rep* 2018;17(4):5409–5415; doi: 10.3892/mmr.2018.8528
33. Gajdzis M, Theocharis S, Gajdzis P, et al. Ephrin receptors (Eph): EphA1, EphA5, and EphA7 expression in uveal melanoma-associations with clinical parameters and patient survival. *Life (Basel)* 2020;10(10):225; doi: 10.3390/life10100225
34. Satofuka HSH, Tanaka T, Li G, et al. An Anti-Human EphA2 Monoclonal Antibody Ea2Mab-7 Shows High Sensitivity for Flow Cytometry, Western Blot, and Immunohistochemical Analyses. Preprint 2024; doi: 10.20944/preprints202410.0572.v1
35. Ubukata R, Suzuki H, Hirose M, et al. Establishment of a highly sensitive and specific anti-EphB2 monoclonal antibody (Eb2Mab-12) for flow cytometry. *MI* 2025;0(0):5728; doi: 10.36922/mi.5728

36. Nanamiya R, Suzuki H, Kaneko MK, et al. Development of an Anti-EphB4 Monoclonal Antibody for Multiple Applications Against Breast Cancers. *Monoclon Antib Immunodiagn Immunother* 2023;42(5):166–177; doi: 10.1089/mab.2023.0015
37. Tanaka T, Kaneko Y, Yamamoto H, et al. Development of a novel anti-erythropoietin-producing hepatocellular receptor B6 monoclonal antibody Eb6Mab-3 for flow cytometry. *Biochem Biophys Rep* 2025;41:101960; doi: 10.1016/j.bbrep.2025.101960
38. Fujii Y, Kaneko MK, Kato Y. MAP Tag: A Novel Tagging System for Protein Purification and Detection. *Monoclon Antib Immunodiagn Immunother* 2016;35(6):293–299; doi: 10.1089/mab.2016.0039
39. Fujii Y, Kaneko M, Neyazaki M, et al. PA tag: A versatile protein tagging system using a super high affinity antibody against a dodecapeptide derived from human podoplanin. *Protein Expr Purif* 2014;95:240–247; doi: 10.1016/j.pep.2014.01.009
40. Ikota H, Nobusawa S, Arai H, et al. Evaluation of IDH1 status in diffusely infiltrating gliomas by immunohistochemistry using anti-mutant and wild type IDH1 antibodies. *Brain Tumor Pathol* 2015;32(4):237–244; doi: 10.1007/s10014-015-0222-8
41. Chu M, Zhang C. Inhibition of angiogenesis by leflunomide via targeting the soluble ephrin-A1/EphA2 system in bladder cancer. *Sci Rep* 2018;8(1):1539; doi: 10.1038/s41598-018-19788-y
42. Herath NI, Boyd AW. The role of Eph receptors and ephrin ligands in colorectal cancer. *Int J Cancer* 2010;126(9):2003–2011; doi: 10.1002/ijc.25147
43. Wang X, Liu Y, Cao G, et al. Expression of the EphA1 protein is associated with Fuhrman nuclear grade in clear cell renal cell carcinomas. *Int J Clin Exp Pathol* 2015;8(6):6821–6827.
44. Arimori T, Mihara E, Suzuki H, et al. Locally misfolded HER2 expressed on cancer cells is a promising target for development of cancer-specific antibodies. *Structure* 2024;32(5):536–549.e535; doi: 10.1016/j.str.2024.02.007
45. Tanaka T, Suzuki H, Ohishi T, et al. A cancer-specific anti-podoplanin monoclonal antibody, PMab-117-mG(2a) exerts antitumor activities in human tumor xenograft models. *Cells* 2024;13(22):1833; doi: 10.3390/cells13221833
46. Liang Z, Wang X, Dong K, et al. Expression pattern and prognostic value of EPHA/EFNA in breast cancer by bioinformatics analysis: Revealing its importance in chemotherapy. *Biomed Res Int* 2021;2021:5575704; doi: 10.1155/2021/5575704
47. Phan NN, Liu S, Wang CY, et al. Overexpressed gene signature of EPH receptor A/B family in cancer patients-comprehensive analyses from the public high-throughput database. *Int J Clin Exp Pathol* 2020;13(5):1220–1242.
48. Pasquale EB. Eph receptors and ephrins in cancer progression. *Nat Rev Cancer* 2024;24(1):5–27; doi: 10.1038/s41568-023-00634-x
49. McCarron JK, Stringer BW, Day BW, et al. Ephrin expression and function in cancer. *Future Oncol* 2010;6(1):165–176; doi: 10.2217/fon.09.146
50. Janes PW, Vail ME, Gan HK, et al. Antibody targeting of Eph receptors in cancer. *Pharmaceuticals (Basel)* 2020;13(5):88; doi: 10.3390/ph13050088
51. Masters CL, Bateman R, Blennow K, et al. Alzheimer's disease. *Nat Rev Dis Primers* 2015;1:15056; doi: 10.1038/nrdp.2015.56
52. Bellenguez C, Grenier-Boley B, Lambert JC. Genetics of Alzheimer's disease: Where we are, and where we are going. *Curr Opin Neurobiol* 2020;61:40–48; doi: 10.1016/j.conb.2019.11.024
53. Li Y, Laws SM, Miles LA, et al. Genomics of Alzheimer's disease implicates the innate and adaptive immune systems. *Cell Mol Life Sci* 2021;78(23):7397–7426; doi: 10.1007/s00018-021-03986-5
54. Lambert JC, Heath S, Even G, et al.; European Alzheimer's Disease Initiative Investigators. Genome-wide association study identifies variants at CLU and CR1 associated with Alzheimer's disease. *Nat Genet* 2009;41(10):1094–1099; doi: 10.1038/ng.439
55. Owens HA, Thorburn LE, Walsby E, et al. Alzheimer's disease-associated P460L variant of EphA1 dysregulates receptor activity and blood-brain barrier function. *Alzheimers Dement* 2024;20(3):2016–2033; doi: 10.1002/alz.13603
56. Buhl E, Kim YA, Parsons T, et al. Effects of Eph/ephrin signalling and human Alzheimer's disease-associated EphA1 on *Drosophila* behaviour and neurophysiology. *Neurobiol Dis* 2022;170:105752; doi: 10.1016/j.nbd.2022.105752
57. Sahoo AR, Buck M. Structural and functional insights into the transmembrane domain association of Eph receptors. *Int J Mol Sci* 2021;22(16):8593; doi: 10.3390/ijms22168593
58. Chen Y, Fu AK, Ip NY. Bidirectional signaling of ErbB and Eph receptors at synapses. *Neuron Glia Biol* 2008;4(3):211–221; doi: 10.1017/s1740925x09990287
59. Ma J, Wang Z, Chen S, et al. EphA1 activation induces neuropathological changes in a mouse model of parkinson's disease through the CXCL12/CXCR4 signaling pathway. *Mol Neurobiol* 2021;58(3):913–925; doi: 10.1007/s12035-020-02122-x
60. Yao M, Miller GW, Vardarajan BN, et al. Deciphering proteins in Alzheimer's disease: A new Mendelian randomization method integrated with AlphaFold3 for 3D structure prediction. *Cell Genom* 2024;4(12):100700; doi: 10.1016/j.xgen.2024.100700
61. Kato Y, Kaneko MK. A cancer-specific monoclonal antibody recognizes the aberrantly glycosylated podoplanin. *Sci Rep* 2014;4:5924; doi: 10.1038/srep05924

Address correspondence to:

Yukinari Kato

Department of Antibody Drug Development
Tohoku University Graduate School of Medicine
2-1, Seiryomachi, Aoba-ku, Sendai
Miyagi 980-8575
Japan

E-mail: yukinari.kato.e6@tohoku.ac.jp

Received: April 22, 2025

Accepted: June 4, 2025